



Patient Instructions for Blepharoplasty/Oculoplastic Surgeries (Eyelid Surgery)

Preparations for Surgery:

For two weeks before and after surgery **AVOID** aspirin or products containing aspirin. Also avoid ibuprofen, naproxen, Vitamin E, and herbal supplements as these products make the blood more difficult to clot. Dr. Barthelow will provide you with a complete list of medications to avoid. You may, however, take Tylenol.

If you smoke, Dr. Barthelow urges you to quit at ***least two weeks prior to surgery.***

Dr. Barthelow will prescribe antibiotics, pain medication, and nausea medication for you during your final preoperative visit. We recommend filling these prescriptions before the day of surgery for your comfort and convenience.

Contact lenses: Remove contact lenses prior to surgery and do not wear them for one week after surgery. Dr. Barthelow will let you know when it is safe to resume wearing your lenses.

Immediately Following Surgery:

You should plan to be off work for up to five to seven days, although this can vary from person to person.

After surgery you will rest quietly for the first 48 hours. You will be able to walk around the house and perform light daily activities; however, during this time, it is not at all unusual for you to feel some pain, soreness and pressure around the eyelids. This will gradually subside and Dr. Barthelow will give you pain medication to relieve it. You must take the entire prescription of antibiotics.

Be sure to keep your head elevated at all times for one week after surgery. You will need to sleep on two or three pillows in order to keep your head elevated. This is the most important factor that will improve the swelling. Do not bend over, tie your shoes, perform any lifting, and avoid any other activity where your face is below the level of your heart.

You will also apply cold compresses across the eyes every four hours for the first 24 hours following surgery. Take a clean washcloth, soak it in ice water, wring it out and apply across the eyes. This is very soothing and with help greatly with the swelling. If you experience a significant change in vision or a large increase in pain call Dr. Barthelow immediately.

You will be allowed to shower two days after surgery. When taking a shower, you may wash your hair normally with shampoo. Do not let the water beat directly down on your eyelids, but just let the water wash over the incisions. Pat the incisions dry and then apply a small amount of antibiotic cream recommended by Dr. Barthelow. This daily routine will help keep the incisions clean, and will promote wound healing. Do not submerge yourself in a bath, swimming pool, or whirlpool for two weeks.

A few patients react to the anesthetic after surgery with nausea and vomiting. This usually lasts less than 24 hours and should be treated with lots of fluids. Dr. Barthelow will prescribe nausea medicine for during your preoperative visit.

You may use your eyes for reading or TV viewing as frequently as you wish. Dr. Barthelow will prescribe antibiotic ointment. You will apply this twice per day until the sutures are removed. Dr. Barthelow will remove the sutures five to seven days after surgery.

The maximum swelling occurs at about three days and then begins to dramatically improve. Mild bruising typically resolves within 14 days. Dr. Barthelow recommends you rest at home for two days after surgery. You may then go out with sunglasses covering the incisions. You will be "socially presentable" in 10 to 14 days.

Eye Lubrication: Keep your eyes lubricated for the first several days after surgery. The best lubricant is artificial tears. The drops should be used every four hours and whenever there are symptoms of dry eyes. Symptoms after eyelid surgery (e.g., itching, irritation, stinging, redness, pain, even tearing) are almost always due to dryness of the surface of the eye. There is no such thing as too much eye lubrication, so use the drops and ointment liberally.

Additional Post-Operative Instructions:

- You may resume wearing make-up 48 hours after the sutures are removed.
- No heavy exercise or lifting for three weeks following surgery. This will allow the wounds to heal without tension.
- Dr. Barthelow encourages walking immediately after surgery. This activity will greatly minimize the risk of deep clots in your leg veins.
- After surgery your eyelids will be swollen. Sometimes the swelling is significant and you will be unable to open your eyes wide for two or three days.
- Any apparent redness in the whites of the eyeball is only a form of bruising, and will subside during the early healing process.
- It is not abnormal to feel slight itching and tightness of the eyelids during the early healing period.

Sexual Activity: Refrain from sexual activity for one week.

Driving: Do not drive an automobile for one week after surgery.

Please notify Dr. Barthelow if:

- If your vision changes suddenly
- If you experience a significant increase in pain in or around the eye after surgery
- If you develop a temperature above 101.5° F
- If you develop redness (like a sunburn) around your eyes

If you need help or have any questions feel free to call Dr. Barthelow at (530) 891-1900 with your concerns. Dr. Barthelow is on call 24 hours per day, seven days per week and has an answering service to forward calls.

The quality of your cosmetic enhancement may be compromised if you fail to return for any scheduled post-op visits, or follow the pre- and post-operative instructions. Don't hesitate to report any unusual or concerning changes.